arXiv:1611.05965v1 [math.FA] 18 Nov 2016

ANOTHER CHARACTERIZATIONS OF MUCKENHOUPT A_p CLASS

DINGHUAI WANG AND JIANG ZHOU*

Abstract. This manuscript addresses Muckenhoupt A_p weight theory in connection to Morrey and BMO spaces. It is proved that ω belongs to Muckenhoupt A_p class, if and only if Hardy-Littlewood maximal function M is bounded from weighted Lebesgue spaces $L^p(\omega)$ to weighted Morrey spaces $M^p_q(\omega)$ for $1 < q < p < \infty$. As a corollary, if M is (weak) bounded on $M^p_q(\omega)$, then $\omega \in A_p$. The A_p condition also characterizes the boundedness of the Riesz transform R_j and convolution operators T_ϵ on weighted Morrey spaces. Finally, we show that $\omega \in A_p$ if and only if $\omega \in \mathrm{BMO}^{p'}(\omega)$ for $1 \le p < \infty$ and $1/p + 1/p' = 1$.

1. Introduction

For $1 < p < \infty$ and a nonnegative locally integrable function ω on $\mathbb{R}^n$, ω is in the Muckenhoupt A_p class if it satisfies the condition

$$[\omega]_{A_p} := \sup_Q \left(\frac{1}{|Q|}\int_Q \omega(x)dx\right)\left(\frac{1}{|Q|}\int_Q \omega(x)^{-\frac{1}{p-1}}dx\right)^{p-1} < \infty.$$

And a weight function ω belongs to the class A_1 if there exists $C > 0$ such that for every cube Q,

$$\frac{1}{|Q|}\int_Q \omega(x)dx \le C \operatorname*{ess\,inf}_{x\in Q} \omega(x),$$

and the infimum of C is denoted by $[\omega]_{A_1}$. A weight ω is called an A_∞ weight if

$$[\omega]_{A_\infty} := \sup_Q \left(\frac{1}{|Q|}\int_Q \omega(x)dx\right)\exp\left(\frac{1}{|Q|}\int_Q \log \omega(x)^{-1}dx\right) < \infty.$$

In fact, $A_\infty = \bigcup_{1\le p<\infty} A_p$.

Weighted inequalities arise naturally in Fourier analysis, but their use is best justified by the variety of applications in which they appear. For example, the theory of weights

2010 Mathematics Subject Classification. 42B20; 42B25.

Key words and phrases. Characterization, Hardy-Littlewood maximal function, Muckenhoupt A_p class, Weighted Morrey spaces, Weighted BMO space.

The research was supported by National Natural Science Foundation of China (Grant No.11261055).

* Corresponding author, Email: zhoujiangshuxue@126.com. Telephone:+8613009688600.

plays an important role in the study of boundary value problems for Laplace's equation on Lipschitz domains. Other applications of weighted inequalities include vector-valued inequalities, extrapolation of operators and applications to certain classes of integral equation and nonlinear partial differential equation. There are a number of classical results demonstrate that the Muckenhoupt A_p classes are the right collections of weights to do harmonic analysis on weighted spaces. The main results along these lines are the equivalence between the $\omega \in A_p$ condition and the $L^p(\omega)$ boundedness (or weak boundedness) of maximal operator and singular integral operators.

A well known result of Muckenhoupt [13] showed that the Hardy-Littlewood maximal function

$$Mf(x) = \sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y)|dy$$

is (weak) bounded on weighted Lebesgue spaces $L^p(\omega)$ if and only if $\omega \in A_p$ for $1 < p < \infty$ (for the case $n = 1$). Hunt, Muckenhoupt and Wheeden [9] proved that the A_p condition also characterizes the $L^p(\omega)$ boundedness of the Hilbert transform

$$Hf(x) = \frac{1}{\pi} \text{p.v.} \int_{\mathbb{R}} \frac{f(y)}{x-y} dy.$$

Later, Coifman and Fefferman [6] extended the A_p theory to the case $n \geq 1$ and general Calderón-Zygmund operators, they also proved that A_p weights satisfy the crucial reverse Hölder condition.

In 2009, Komori and Shirai [10] introduced the weighted Morrey spaces. Let $0 < q < p < \infty$, ω be a weight and $\omega(Q) := \int_Q \omega(x)dx$. Then a weighted Morrey space is defined by

$$M^p_q(\omega) = \Big\{ f \in L^q_{loc}(\omega) : \|f\|_{M^p_q(\omega)} := \sup_Q \frac{1}{\omega(Q)^{1/q-1/p}} \Big(\int_Q |f(x)|^q \omega(x)dx \Big)^{1/q} < \infty \Big\},$$

and a weighted weak Morrey space is defined by

$$WM^p_q(\omega) = \Big\{ f \in L^q_{loc}(\omega) : \|f\|_{WM^p_q(\omega)} < \infty \Big\},$$

where

$$\|f\|_{WM^p_q(\omega)} := \sup_Q \frac{1}{\omega(Q)^{1/q-1/p}} \sup_{\lambda>0} \lambda \Big(\int_{\{x \in Q : |f(x)| > \lambda\}} \omega(x)dx \Big)^{1/q}.$$

They proved that if $\omega \in A_p$, then M is bounded on $M^p_q(\omega)$. An interesting question is raised. Is ω in A_p if M is bounded on $M^p_q(\omega)$ for $1 < q < p < \infty$? We will give an affirmative answer as follows.

Theorem 1.1. *Let $1 < q < p < \infty$. The following statements are equivalent:*

(1) $\omega \in A_p$;
(2) M *is a bounded operator from* $L^p(\omega)$ *to* $L^{p,\infty}(\omega)$;
(3) M *is a bounded operator from* $L^p(\omega)$ *to* $M^p_q(\omega)$;
(4) M *is a bounded operator from* $L^p(\omega)$ *to* $WM^p_q(\omega)$;
(5) M *is a bounded operator from* $M^p_q(\omega)$ *to* $M^p_q(\omega)$;
(6) M *is a bounded operator from* $M^p_q(\omega)$ *to* $WM^p_q(\omega)$.

Remark 1.1. *It should be point out that the main contribution of this paper in Theorem 1.1 is* (2) $\Rightarrow$ (3) *and* (4) $\Rightarrow$ (1)*, and other implications have been showed in* [6] *and* [10] *or follow from the trivial embedding properties.*

For the case $p = 1$, we have

Theorem 1.2. *Let $0 < q < 1$. The following statements are equivalent:*

(1) $\omega \in A_1$;
(2) M *is a bounded operator from* $L^1(\omega)$ *to* $L^{1,\infty}(\omega)$;
(3) M *is a bounded operator from* $L^1(\omega)$ *to* $M^1_q(\omega)$;
(4) M *is a bounded operator from* $L^1(\omega)$ *to* $WM^1_q(\omega)$.

For the nth dimensional case, the A_p condition also characterizes the $L^p(\omega)$ boundedness of the Riesz transform

$$R_j f(x) = c_n \text{p.v.} \int_{\mathbb{R}^n} \frac{x_j - y_j}{|x-y|^{n+1}} f(y)dy, \quad j = 1, 2, \cdots, n,$$

where $c_n = \Gamma(\frac{n+1}{2})/\pi^{\frac{n+1}{2}}$. In this paper, we will show that the A_p condition is also necessary for the boundedness of the Riesz transforms on weighted Morrey spaces.

Theorem 1.3. *Let $1 < q < p < \infty$. The following statements are equivalent:*

(1) $\omega \in A_p$;
(2) R_j *is a bounded operator from* $L^p(\omega)$ *to* $L^p(\omega)$ *for all* $j = 1, 2, \cdots, n$;
(3) R_j *is a bounded operator from* $L^p(\omega)$ *to* $M^p_q(\omega)$ *for all* $j = 1, 2, \cdots, n$;
(4) R_j *is a bounded operator from* $L^p(\omega)$ *to* $WM^p_q(\omega)$ *for all* $j = 1, 2, \cdots, n$;
(5) R_j *is a bounded operator from* $M^p_q(\omega)$ *to* $M^p_q(\omega)$ *for all* $j = 1, 2, \cdots, n$;
(6) R_j *is a bounded operator from* $M^p_q(\omega)$ *to* $WM^p_q(\omega)$ *for all* $j = 1, 2, \cdots, n$.

For the case $p = 1$, we have

Theorem 1.4. *Let $0 < q < 1$. The following statements are equivalent:*

(1) $\omega \in A_1$;
(2) R_j *is a bounded operator from* $L^1(\omega)$ *to* $L^{1,\infty}(\omega)$ *for all* $j = 1, 2, \cdots, n$;
(3) R_j *is a bounded operator from* $L^1(\omega)$ *to* $M^1_q(\omega)$ *for all* $j = 1, 2, \cdots, n$;
(4) R_j *is a bounded operator from* $L^1(\omega)$ *to* $WM^1_q(\omega)$ *for all* $j = 1, 2, \cdots, n$.

In [17, page 198], Stein showed that the convolution operators T_ϵ is bounded on weighted Lebesgue spaces if and only if $\omega \in A_p$. The convolution operators T_ϵ are defined by

$$T_\epsilon f = f * \Phi_\epsilon,$$

where Φ is nonnegative, radial, and (radially) decreasing, with $\int \Phi(x)dx = 1$, and we define $\Phi_\epsilon(x) = \epsilon^{-n}\Phi(x/\epsilon)$. We designate the class of such Φ by $\mathcal{R}$. Notice that if Φ is in $\mathcal{R}$, then so is Φ_ϵ. It is useful to recall that

$$Mf(x) = \sup_{\Phi\in\mathcal{R}} |f| * \Phi(x).$$

In fact, if $B_\epsilon = \{x : |x| < \epsilon\}$ then $|B_\epsilon|^{-1}\chi_{B_\epsilon} \in \mathcal{R}$, and the supremum over these elements of $\mathcal{R}$ is, by definition, equal to $Mf(x)$. For the other direction, we need only recall that any element of $\mathcal{R}$ is a limit of weighted averages of the $|B_\epsilon|^{-1}\chi_{B_\epsilon}$ (see [17, Chapter 2, §2.1]).

Theorem 1.5. *Let $1 < q < p < \infty$. The following statements are equivalent:*

(1) $\omega \in A_p$;
(2) T_ϵ *is a bounded operator from* $L^p(\omega)$ *to* $L^p(\omega)$;
(3) T_ϵ *is a bounded operator from* $L^p(\omega)$ *to* $L^{p,\infty}(\omega)$;
(4) T_ϵ *is a bounded operator from* $L^p(\omega)$ *to* $M^p_q(\omega)$;
(5) T_ϵ *is a bounded operator from* $L^p(\omega)$ *to* $WM^p_q(\omega)$;
(6) T_ϵ *is a bounded operator from* $M^p_q(\omega)$ *to* $M^p_q(\omega)$;
(7) T_ϵ *is a bounded operator from* $M^p_q(\omega)$ *to* $WM^p_q(\omega)$.

Finally, we also consider Muckenhoupt weight theory related to weighted BMO spaces. Weighted BMO spaces play a fundamental role in many fields of mathematics such as harmonic analysis and partial differential equations; see [3], [4], [8] and [12]. Let us introduce the weighted BMO spaces.

Let $1 \le p < \infty$. Given a a nonnegative locally integrable function ω, the weighted BMO space $\mathrm{BMO}^p(\omega)$ is defined be the set of all functions $f \in L^1_{\mathrm{loc}}(\mathbb{R}^n)$ such that

$$\|f\|_{\mathrm{BMO}^p(w)} := \sup_Q \left(\frac{1}{w(Q)}\int_Q |f(y) - f_Q|^p \omega(y)^{1-p}dy\right)^{1/p} < \infty,$$

where the supremum is taken over all cubes $Q \subset \mathbb{R}^n$. We write $\mathrm{BMO}^1(\omega) = \mathrm{BMO}(\omega)$ simple.

Remark 1.2. (i) For $1 \leq p < \infty$ and $\omega \in A_1$, García-Cuerva [8] proved that $\mathrm{BMO}(\omega) = \mathrm{BMO}^p(\omega)$ with equivalence of the corresponding norms.

(ii) For the case $p = \infty$, the weighted BMO space can be defined as

$$\|f\|_{\mathrm{BMO}^\infty(\omega)} := \operatorname*{ess\,sup}_{x \in Q} \frac{|f(x) - f_Q|}{\omega(x)}.$$

Now we state the result of the characterization of Muckenhoupt A_p class via weighted BMO spaces.

Theorem 1.6. *Let $1 \leq p < \infty$ and $\frac{1}{p} + \frac{1}{p'} = 1$. Given a nonnegative locally integrable function ω, $\omega \in A_p$ if and only if $\omega \in \mathrm{BMO}^{p'}(\omega)$. Moreover,*

$$[\omega]_{A_p}^{1/p} \leq \|\omega\|_{\mathrm{BMO}^{p'}(\omega)} + 1 \leq 3[\omega]_{A_p}^{1/p}.$$

For the case $p = \infty$, we have the following result.

Theorem 1.7. *Given a nonnegative locally integrable function $\omega \in A_\infty$, then $\omega \in \mathrm{BMO}(\omega)$.*

Though the proof of Theorem 1.6 is fairly straightforward, this result seems interesting. It appears to provide some insight into the growth allowed for Muckenhoupt weights. As an application, the operator norms of weighted Hardy-Littlewood average operator U_ψ on weighted BMO spaces are also obtained.

Theorem 1.8. *Let $\psi : [0,1] \to [0,\infty)$ be a function, $p \in [1,\infty]$, $\alpha \in \mathbb{R}$, $\omega \in A_{p'}$ and $\omega(tx) = t^\alpha \omega(x)$ for all $t \in (0,\infty)$. Then $U_\psi : \mathrm{BMO}^p(\omega) \to \mathrm{BMO}^p(\omega)$ exists as a bounded operator if and only if*

$$\int_0^1 t^\alpha \psi(t) dt < \infty. \tag{1.1}$$

Moreover, when (1.1) holds, the operator norm of U_ψ on $\mathrm{BMO}^p(\omega)$ is given by

$$\|U_\psi\|_{\mathrm{BMO}^p(\omega) \to \mathrm{BMO}^p(\omega)} = \int_0^1 t^\alpha \psi(t) dt.$$

Remark 1.3. (i) *If $\omega(x) = |x|^\alpha$, the condition $\omega(tx) = t^\alpha \omega(x)$ holds.*

(ii) *The norm from $\mathrm{BMO}^p(\omega)$ to itself of the operator U_ψ is independent of p.*

The definition of weighted Hardy-Littlewood average operator U_ψ as follows.

Let $\psi : [0,1] \to [0,\infty)$ be a function. For a measurable complex valued function f on $\mathbb{R}^n$, the weighted Hardy-Littlewood average operator U_ψ is defined as

$$(U_\psi f)(x) = \int_0^1 f(tx)\psi(t)dt.$$

It was first defined by Carton-Lebrun and Fosset in [5] and they showed that if $t^{1-n}\psi(t)$ is bounded on $[0,1]$ then U_ψ is bounded on $\mathrm{BMO}(\mathbb{R}^n)$. In 2001, Xiao [18] obtained that U_ψ is bounded on $L^p(\mathbb{R}^n)$, $1 \le p \le \infty$, if and only if

$$\int_0^1 t^{-n/p}\psi(t)dt < \infty. \tag{1.2}$$

Meanwhile, when (1.2) holds,

$$\|U_\psi\|_{L^p(\mathbb{R}^n)\to L^p(\mathbb{R}^n)} = \int_0^1 t^{-n/p}\psi(t)dt.$$

Xiao also showed that U_ψ is bounded on $\mathrm{BMO}(\mathbb{R}^n)$ if and only if

$$\int_0^1 \psi(t)dt < \infty. \tag{1.3}$$

And when (1.3) holds, the precise norm of U_ψ on $\mathrm{BMO}(\mathbb{R}^n)$ is given by

$$\|U_\psi\|_{\mathrm{BMO}(\mathbb{R}^n)\to\mathrm{BMO}(\mathbb{R}^n)} = \int_0^1 \psi(t)dt.$$

This paper is organized as follows. In Section 2, we prove that A_p weights give a characterization of weighted Morrey spaces $M^p_q(\omega)$ boundedness for the Hardy-Littlewood maximal operator (see Theorem 1.1 and Theorem 1.2), Riesz transform (see Theorem 1.3 and Theorem 1.4) and convolution operators T_ϵ (see Theorem 1.5). In Section 3, we address A_p weight theory in connection to weighted BMO spaces and its application (see Theorem 1.6, Theorem 1.7 and Theorem 1.8).

Throughout this paper, the letter C denotes constants which are independent of main variables and may change from one occurrence to another.

2. Proof of Theorem 1.1 $\sim$ Theorem 1.5

First of all, we compare with weighted Morrey spaces and weighted weak Morrey spaces. It is clear that $M^p_r(\omega)$ is contained in $WM^p_q(\omega)$ and $\|\cdot\|_{WM^p_q(\omega)} \le \|\cdot\|_{M^p_q(\omega)} \le \|\cdot\|_{M^p_r(\omega)}$ if $1 < q \le r < p < \infty$. However, for $1 < r < q \le p < \infty$, one has the reverse inequality as follows.

Proposition 2.1. *Let $0 < q \le p < \infty$ and ω be a nonnegative locally integral function on $\mathbb{R}^n$.*

(1) *If $f \in WM_q^p(\omega)$, then for all $r \in (0,q)$ and all cube Q, there exists a constant $C > 0$ such that*

$$\int_Q |f(x)|^r \omega(x)dx \le C\omega(Q)^{1-r/p}\|f\|^r_{WM_q^p(\omega)}.$$

(2) *If there exists $r \in (0,q)$ and constant $C > 0$ such that*

$$\int_\Omega |f(x)|^r \omega(x)dx \le C\omega(\Omega)^{1-r/p} \tag{2.1}$$

for all set Ω with $\omega(\Omega) < \infty$, then $f \in WM_q^p(\omega)$.

Proof. Let $f \in WM_q^p(\omega)$. For any $\lambda > 0$

$$\lambda\Big(\int_{\{x\in Q:|f(x)|>\lambda\}} \omega(x)dx\Big)^{1/q} \le \|f\|_{WM_q^p(\omega)}\omega(Q)^{1/q-1/p}.$$

Choose

$$N = \|f\|_{WM_q^p(\omega)}\big(\frac{r}{q-r}\big)^{1/q}\omega(Q)^{-1/p},$$

for any $0 < r < q$,

$$\begin{aligned}\int_Q |f(x)|^r\omega(x)dx &= r\int_0^\infty \lambda^{r-1}\omega\big(\{x\in Q:|f(x)|>\lambda\}\big)d\lambda\\ &\le r\int_0^N \lambda^{r-1}\omega(Q)d\lambda + r\int_N^\infty \lambda^{r-1}\frac{\|f\|^q_{WM_q^p(\omega)}}{\lambda^q}\omega(Q)^{1-q/p}d\lambda\\ &= N^r\omega(Q) + \|f\|^p_{WM_q^p(\omega)}\frac{r}{q-r}N^{r-q}\omega(Q)^{1-q/p},\end{aligned}$$

which implies

$$\Big(\int_Q |f(x)|^r\omega(x)dx\Big)^{1/r} \le 2\|f\|_{WM_q^p(\omega)}\big(\frac{r}{q-r}\big)^{1/q}\omega(Q)^{1/r-1/p};$$

that is,

$$\int_Q |f(x)|^r\omega(x)dx \le C\omega(Q)^{1-r/p}\|f\|^r_{WM_q^p(\omega)}.$$

Proof of (2). For any $\lambda > 0$, take $E = \{x \in \mathbb{R}^n : |f(x)| > \lambda\}$, then $\omega(E) < \infty$. Otherwise, there is a sequence $\{E_k\}$ of measurable sets such that $E_k \subset E$ and $\omega(E_k) = k$ for $k = 0, 1, 2, \cdots$. Thus for every k,

$$\lambda^r k = \lambda^r\omega(E_k) \le \int_{E_k} |f(x)|^r\omega(x)dx \le C\omega(E_k)^{1-r/p} = Ck^{1-r/p}.$$

However, it is not true.

Thus, for any cube $Q \subset \mathbb{R}^n$, take $\Omega = \{x \in Q : |f(x)| > \lambda\}$(for $p = q$, take $\Omega = E$), then $\omega(\Omega) \le \omega(E) < \infty$. By (2.1), we have

$$\lambda^r \omega(\Omega) \le \int_\Omega |f(x)|^r \omega(x)dx \le C\omega(\Omega)^{1-r/p}.$$

It follows that

$$\lambda\omega(Q)^{1/p-1/q}\omega(\Omega)^{1/q} \le \lambda\omega(\Omega)^{1/p} \le C.$$

Hence

$$\sup_Q \frac{1}{\omega(Q)^{1/q-1/p}} \sup_{\lambda>0} \lambda\Big(\int_{\{x\in Q:|f(x)|>\lambda\}} \omega(x)dx\Big)^{1/q} \le C,$$

then $f \in WM^p_q(\omega)$. □

Remark 2.1. In fact, for $p = q$, we can replace "for all cube Q" in Proposition 2.1 (1) with "for all set Ω with $\omega(\Omega) < \infty$". For the proof similar arguments are applied with necessary modifications.

As a corollary of Proposition 2.1, we get

Corollary 2.1. *Let $0 < r < q < p < \infty$ and ω be a nonnegative locally integral function on $\mathbb{R}^n$. Weighted Morrey space $M^p_r(\omega)$ is contained in $WM^p_q(\omega)$ and*

$$\|\cdot\|_{M^p_r(\omega)} \le C\|\cdot\|_{WM^p_q(\omega)} \le C\|\cdot\|_{M^p_q(\omega)} \le C\|\cdot\|_{WM^p_p(\omega)} = C\|\cdot\|_{L^{p,\infty}(\omega)}.$$

Proof of Theorem 1.1. The equivalence of (1) and (2) was proved in [6]. By Corollary 2.1, it is obvious that (2) implies (3), and (3) implies (4) follows from the trivial embedding properties.

(4) ⇒ (1): Let Q be any cube and assuming for a moment that $\int_Q \omega(x)^{1-p'}dx < \infty$. We take $f = \omega^{1-p'}\chi_Q$. For any $0 < \lambda < \omega^{1-p'}(Q)/|Q|$, we obtain that for any $x \in Q$, $Mf(x) > \lambda$. Then

$$\begin{aligned}
\frac{\lambda\omega(Q)^{1/p}}{|Q|^{1/p}} &= \frac{\lambda}{|Q|^{1/p}} \frac{1}{\omega(Q)^{1/q-1/p}} \Big(\int_Q \omega(x)dx\Big)^{1/q} \\
&= \frac{1}{|Q|^{1/p}} \frac{\lambda}{\omega(Q)^{1/q-1/p}} \Big(\int_{\{x\in Q:\ Mf(x)>\lambda\}} \omega(x)dx\Big)^{1/q} \\
&\le \frac{C}{|Q|^{1/p}} \Big(\int_{\mathbb{R}^n} [\omega(x)^{1-p'}\chi_Q(x)]^p \omega(x)dx\Big)^{1/p} \\
&= \frac{C}{|Q|^{1/p}} \Big(\int_Q \omega(x)^{1-p'}dx\Big)^{1/p}.
\end{aligned}$$

Hence by arbitrariness of λ, we get

$$\frac{1}{|Q|}\int_Q \omega(x)dx \le C\Big(\frac{1}{|Q|}\int_Q \omega(x)^{1-p'}dx\Big)^{1-p};$$

that is,

$$\Big(\frac{1}{|Q|}\int_Q \omega(x)dx\Big)\Big(\frac{1}{|Q|}\int_Q \omega(x)^{1-p'}dx\Big)^{p-1} \le C. \tag{2.2}$$

If $\omega(x)^{1-p'}$ is not locally integrable, then we take $f=(\omega+\epsilon)^{1-p'}\chi_Q$ and $0<\lambda<(\omega+\epsilon)^{1-p'}(Q)/|Q|$ to get

$$\Big(\frac{1}{|Q|}\int_Q \omega(x)dx\Big)\Big(\frac{1}{|Q|}\int_Q (\omega(x)+\epsilon)^{-p'}\omega(x)dx\Big)^{p-1} \le C,$$

for all $\epsilon>0$, from which we can still deduce (2.2) by letting $\epsilon\to 0$. So $\omega\in A_p$.

In addition, (1) $\Rightarrow$ (5) was proved in [10], and it is obvious that (5) $\Rightarrow$ (6). Since $\|\cdot\|_{M^p_q(\omega)} \le \|\cdot\|_{L^p(\omega)}$ for any weight function ω and $1<q<p<\infty$, we have (6) $\Rightarrow$ (4). Finally, (4) $\Rightarrow$ (1) can be found above. □

Proof of Theorem 1.2. We only give the proof of (4) $\Rightarrow$ (1), and other implications have been showed in [6], Corollary 2.1 or follow from the trivial embedding properties.

(4) $\Rightarrow$ (1): Let Q be any cube and $Q_1\subset Q$. If denote $f=\chi_{Q_1}$, then for any $0<\lambda<\frac{|Q_1|}{|Q|}$, we have

$$Q=\big\{x\in Q: Mf(x)>\lambda\big\}.$$

Applying M is bounded from $L^1(\omega)$ to $WM^1_q(\omega)$, we get

$$\lambda\int_Q \omega(x)dx = \lambda\omega(Q)^{1-1/q}\Big(\int_{\{x\in Q: Mf(x)>\lambda\}}\omega(x)dx\Big)^{1/q} \le C\int_{Q_1}\omega(x)dx.$$

Choose $\lambda=\frac{|Q_1|}{2|Q|}$, then

$$\frac{1}{|Q|}\int_Q \omega(x)dx \le \frac{C}{|Q_1|}\int_{Q_1}\omega(x)dx.$$

It follows from Lebesgue differentiation theorem that

$$\frac{1}{|Q|}\int_Q \omega(x)dx \le C\omega(x) \quad a.e. \quad x\in Q,$$

which proves

$$M\omega(x)\le C\omega(x) \quad a.e. \quad x\in\mathbb{R}^n.$$

Thus $\omega\in A_1$. □

Sharp bounds for the operators norms in terms of the A_p constants of the weights have been investigated as well. Buckley [2] obtained that for $1 < p < \infty$,

$$\|M\|_{L^p(\omega)\to L^p(\omega)} \le C[\omega]_{A_p}^{1/(p-1)}$$

and the power $[\omega]_{A_p}^{1/(p-1)}$ is the best possible. The weak type bound was found and shown to be best possible by Muckenhoupt [13], when the proof is examined closely; that is,

$$\|M\|_{L^p(\omega)\to L^{p,\infty}(\omega)} \le C[\omega]_{A_p}^{1/p} \tag{2.3}$$

and the power $[\omega]_{A_p}^{1/p}$ is the best possible. However, only partial results of singular integrals operators are known. The interest in sharp weighted norm for singular integral operators is motivated in part by applications in partial differential equations. We refer the reader to [1], [7], [11], [14], [15] and [16].

Now, we give the sharp estimate for the boundedness of M from weighted Lebesgue spaces to weighted (weak) Morrey spaces.

Theorem 2.2. *If $1 < q < p < \infty$ and $\omega \in A_p$, then $\|Mf\|_{M_q^p(\omega)} \le C[\omega]_{A_p}^{1/p}\|f\|_{L^p(\omega)}$ and the power $[\omega]_{A_p}^{1/p}$ is best possible.*

Theorem 2.3. *If $1 < q < p < \infty$ and $\omega \in A_p$, then $\|Mf\|_{WM_q^p(\omega)} \le C[\omega]_{A_p}^{1/p}\|f\|_{L^p(\omega)}$ and the power $[\omega]_{A_p}^{1/p}$ is best possible.*

We focus on the proof of Theorem 2.2. For the proof of Theorem 2.3 similar arguments are applied with necessary modifications, we omit the details.

Proof of Theorem 2.2. On the one hand, by Corollary 2.1 and the inequalities (2.3), we have

$$\|Mf\|_{M_q^p(\omega)} \le C\|Mf\|_{L^{p,\infty}(\omega)} \le C[\omega]_{A_p}^{1/p}\|f\|_{L^p(\omega)}.$$

On the other hand, let Q be any cube, take $f = \omega^{1-p'}\chi_Q$ and $\lambda = \frac{\omega^{1-p'}(Q)}{2|Q|}$ (If $\omega(x)^{1-p'}$ is not locally integrable, then we take $f = (\omega+\epsilon)^{-p'/p}\chi_Q$ and $\lambda = \frac{(\omega+\epsilon)^{1-p'}(Q)}{2|Q|}$). Since

$$\{x : x \in Q\} = \{x \in Q : \ Mf(x) > \lambda\},$$

then

$$\begin{aligned}\frac{\lambda\omega(Q)^{1/p}}{|Q|^{1/p}} &= \frac{\lambda}{|Q|^{1/p}}\frac{1}{\omega(Q)^{1/q-1/p}}\Big(\int_Q \omega(x)dx\Big)^{1/q} \\ &= \frac{1}{|Q|^{1/p}}\frac{\lambda}{\omega(Q)^{1/q-1/p}}\Big(\int_{\{x\in Q:\ Mf(x)>\lambda\}} \omega(x)dx\Big)^{1/q}\end{aligned}$$

$$\begin{aligned}
&\leq \frac{1}{|Q|^{1/p}}\frac{1}{\omega(Q)^{1/q-1/p}}\Big(\int_Q |Mf(x)|^q\omega(x)dx\Big)^{1/q}\\
&\leq \frac{\|M\|_{L^p(\omega)\to M^p_q(\omega)}}{|Q|^{1/p}}\Big(\int_{\mathbb{R}^n}[\omega(x)^{1-p'}\chi_Q(x)]^p\omega(x)dx\Big)^{1/p}\\
&= \frac{\|M\|_{L^p(\omega)\to M^p_q(\omega)}}{|Q|^{1/p}}\Big(\int_Q \omega(x)^{1-p'}dx\Big)^{1/p},
\end{aligned}$$

which implies that

$$[\omega]_{A_p} \leq 2^p\|M\|^p_{L^p(\omega)\to M^p_q(\omega)}$$

and Theorem 2.2 follows. □

Similar to Theorem 1.1 and Theorem 1.2, we only prove (4) ⇒ (1) of Theorem 1.3 and Theorem 1.4. In fact, other implications have been showed in [6] and [10] or follow from the trivial embedding properties.

Proof of Theorem 1.3 and Theorem 1.4. (4) ⇒ (1): We prove the nth dimensional case when $n \geq 2$. The one-dimensional case is obtained by a simple adaptation of the following argument.

Let Q be a cube and let f be a nonnegative function on $\mathbb{R}^n$ supported in Q that satisfies $f_Q > 0$. Let Q' be the cube that shares a corner with Q, which has the same length as Q with $x_j \geq y_j$ for all $x \in Q'$ and $y \in Q$. Then for $x \in Q'$ and $y \in Q$, we have $\sum_{j=1}^n x_j - y_j \geq |x-y|$ and $|x-y|^{-n} \geq C|Q|^{-1}$, which implies that for $x \in Q'$,

$$\Big|\sum_{j=1}^n R_jf(x)\Big| = c_n\Big|\sum_{j=1}^n\int_{\mathbb{R}^n}\frac{x_j-y_j}{|x-y|^{n+1}}f(y)dy\Big| \geq C\int_Q\frac{f(y)}{|x-y|^n} \geq Cf_Q.$$

For all $0 < \lambda < Cf_Q$ we have

$$Q' = \big\{x \in Q' : \Big|\sum_{j=1}^n R_jf(x)\Big| > \lambda\big\}.$$

Since the operator $\sum_{j=1}^n R_j$ is bounded from $L^p(\omega)$ to $WM^p_q(\omega)$ for $p \geq 1$, then

$$\omega(Q') \leq \frac{C}{\lambda^p}\int_Q f(x)^p\omega(x)dx,$$

for all $0 < \lambda < Cf_Q$, it proves that

$$f_Q^p \leq \frac{C}{\omega(Q')}\int_Q f(x)^p\omega(x)dx. \tag{2.4}$$

We observe that we can reverse the roles of Q and Q' and obtain

$$g_{Q'}^p \le \frac{C}{\omega(Q)} \int_{Q'} g(x)^p \omega(x) dx, \tag{2.5}$$

for all g supported in Q'. Taking $g = \chi_{Q'}$ in (2.5), which gives that

$$\omega(Q) \le C\omega(Q').$$

Using this estimate and (2.5), we obtain

$$f_Q^p \le \frac{C}{\omega(Q)} \int_Q f(x)^p \omega(x) dx. \tag{2.6}$$

Case 1: $p > 1$. Taking $f = (\omega + \epsilon)^{-p'/p} \chi_Q$ in (2.6), we have

$$\Big(\frac{1}{|Q|}\int_Q \omega(x)dx\Big)\Big(\frac{1}{|Q|}\int_Q (\omega(x)+\epsilon)^{1-p'}dx\Big)^p\Big(\frac{1}{|Q|}\int_Q (\omega(x)+\epsilon)^{-p'}\omega(x)dx\Big)^{-1} \le C.$$

Letting $\epsilon \to 0$ and it follows that $\omega \in A_p$.

Case 2: $p = 1$. For any $Q_1 \subset Q$ and denote $f = \chi_{Q_1}$. By (2.6),

$$\frac{1}{|Q|}\int_Q \omega(x)dx \le \frac{C}{|Q_1|}\int_{Q_1} \omega(x)dx.$$

Applying Lebesgue differentiation theorem that

$$\frac{1}{|Q|}\int_Q \omega(x)dx \le C\omega(x) \quad a.e. \quad x \in Q.$$

Thus $\omega \in A_1$. □

Proof of Theorem 1.5. We need only to show that (5) ⇒ (1). In fact, (1) ⇒ (2) has been proved in [17, page 198] and (1) ⇒ (6) can be obtained by the fact that $|T_\epsilon f| \le Mf$(see [17, page 57]) and the boundedness of M on weighted Morrey spaces, other implications follow from the trivial embedding properties.

(5) ⇒ (1): We first assume that $\Phi_\epsilon = |B_\epsilon|^{-1}\chi_{B_\epsilon}$. Let $B := B(x_0, \delta)$ be any ball and let $f \ge 0$ supported in B. If $x \in B$, then $B \subset R := B(x, 2\delta)$. Thus if $x \in B$ and $\epsilon = 2\delta$, we have

$$\begin{aligned} T_\epsilon f(x) = f * \Phi_\epsilon(x) &= \int_{\mathbb{R}^n} f(y)\Phi_\epsilon(x-y)dy \\ &= \int_{\mathbb{R}^n} f(y)|B_\epsilon|^{-1}\chi_{B_\epsilon}(x-y)dy \\ &= |B_\epsilon|^{-1}\int_{|x-y|<2\delta} f(y)dy \end{aligned}$$

$$\geq \ 2^{-n}\frac{1}{|B|}\int_B f(y)dy = 2^{-n}f_B.$$

For any $0<\lambda<2^{-n}f_B$, by the fact that T_ϵ is a bounded operator from $L^p(\omega)$ to $WM^p_q(\omega)$, we obtain

$$\begin{aligned} C\bigg(\int_{\mathbb{R}^n} f(x)^p\omega(x)dx\bigg)^{1/p} &\geq \omega(B)^{1/p-1/q}\sup_{\lambda>0}\lambda\bigg(\int_{\{x\in B:\ T_\epsilon f(x)>\lambda\}}\omega(x)dx\bigg)^{1/q} \\ &\geq \omega(B)^{1/p}\cdot 2^{-n}f_B. \end{aligned}$$

Taking $f=(\omega+\epsilon)^{-p'/p}\chi_B$, the above inequality implies that

$$\Big(\frac{1}{|Q|}\int_Q\omega(x)dx\Big)\Big(\frac{1}{|Q|}\int_Q(\omega(x)+\epsilon)^{1-p'}dx\Big)^p\Big(\frac{1}{|Q|}\int_Q(\omega(x)+\epsilon)^{-p'}\omega(x)dx\Big)^{-1}\leq C.$$

Letting $\epsilon\to 0$ and it follows that $\omega\in A_p$.

For a general $\Phi\in\mathcal{R}$, there exist some positive constants c_1 and c_2 such that $\Phi(x)\geq c_1\chi_{B_{c_2}}(x)$ (see [17, page 199]), then the same conclusion follows. □

3. Proof of Theorem 1.6 ∼ Theorem 1.8

Below we shall turn to discuss the properties of weighted BMO spaces.

Proposition 3.1. *Let $1\leq p\leq\infty$, $1/p+1/p'=1$ and $\omega\in A_p$. If $f\in \mathrm{BMO}^{p'}(\omega)$, then $|f|\in\mathrm{BMO}^{p'}(\omega)$. Moreover,*

$$\big\||f|\big\|_{\mathrm{BMO}^{p'}(\omega)}\leq(1+[\omega]_{A_p}^{1/p})\|f\|_{\mathrm{BMO}^{p'}(\omega)}.$$

Proof. Let $f\in\mathrm{BMO}^{p'}(\omega)$.

Case 1: $p=1$. For any cube $x\in Q$,

$$\begin{aligned} \big||f(x)|-(|f|)_Q\big| &\leq \big||f(x)|-f_Q\big|+\big|f_Q-(|f|)_Q\big| \\ &\leq \big|f(x)-f_Q\big|+\frac{1}{|Q|}\int_Q|f(y)-f_Q|dy \\ &\leq \big|f(x)-f_Q\big|+M\omega(x)\|f\|_{\mathrm{BMO}^\infty(\omega)}. \end{aligned}$$

It follows from $\omega\in A_1$ that

$$M\omega(x)\leq[\omega]_{A_1}\omega(x)\quad a.e.\quad x\in\mathbb{R}^n.$$

Then

$$\big\||f|\big\|_{\mathrm{BMO}^\infty(\omega)}\leq(1+[\omega]_{A_1})\|f\|_{\mathrm{BMO}^\infty(\omega)}.$$

Case 2: $1 < p < \infty$. Since $\omega \in A_p$, we have

$$\begin{aligned}\big||f(x)| - (|f|)_Q\big| &\le \big||f(x)| - |f_Q|\big| + \big||f_Q| - (|f|)_Q\big| \\ &\le \big|f(x) - f_Q\big| + \frac{1}{|Q|}\int_Q |f(y) - f_Q|dy \\ &\le \big|f(x) - f_Q\big| + \frac{1}{|Q|}\Big(\int_Q |f(y) - f_Q|^{p'}\omega(y)^{1-p'}dy\Big)^{1/p'}\omega(Q)^{1/p} \\ &\le \big|f(x) - f_Q\big| + \frac{\omega(Q)}{|Q|}\|f\|_{\mathrm{BMO}^{p'}(\omega)},\end{aligned}$$

which yields that

$$\begin{aligned}&\left(\frac{1}{\omega(Q)}\int_Q \big||f(x)| - (|f|)_Q\big|^{p'}\omega(x)^{1-p'}dx\right)^{1/p'} \\ &\le \left(\frac{1}{\omega(Q)}\int_Q |f(x) - f_Q|^{p'}\omega(x)^{1-p'}dx\right)^{1/p'} \\ &\quad + \left(\frac{1}{\omega(Q)}\int_Q \Big(\frac{\omega(Q)}{|Q|}\|f\|_{\mathrm{BMO}^p(\omega)}\Big)^{p'}\omega(x)^{1-p'}dx\right)^{1/p'} \\ &\le \|f\|_{\mathrm{BMO}^{p'}(\omega)} + \|f\|_{\mathrm{BMO}^{p'}(\omega)}\left(\Big(\frac{1}{|Q|}\int_Q \omega(x)dx\Big)\Big(\frac{1}{|Q|}\int_Q \omega(x)^{1-p'}dx\Big)^{p-1}\right)^{1/p} \\ &\le (1 + [\omega]_{A_p}^{1/p})\|f\|_{\mathrm{BMO}^{p'}(\omega)}.\end{aligned}$$

Case 3: $p = \infty$. For any cube Q,

$$\begin{aligned}\big||f(x)| - (|f|)_Q\big| &\le \big||f(x)| - |f_Q|\big| + \big||f_Q| - (|f|)_Q\big| \\ &\le \big|f(x) - f_Q\big| + \frac{1}{|Q|}\int_Q |f(y) - f_Q|dy \\ &\le \big|f(x) - f_Q\big| + \frac{\omega(Q)}{|Q|}\|f\|_{\mathrm{BMO}(\omega)}.\end{aligned}$$

It is easy to see that

$$\frac{1}{\omega(Q)}\int_Q \big||f(x)| - (|f|)_Q\big|dx \le 2\|f\|_{\mathrm{BMO}(\omega)}.$$

Combining the estimates above, we conclude that

$$\big\||f|\big\|_{\mathrm{BMO}^{p'}(\omega)} \le (1 + [\omega]_{A_p}^{1/p})\|f\|_{\mathrm{BMO}^{p'}(\omega)}.$$

Thus, we complete the proof of Proposition 3.1. □

Proof of Theorem 1.5. On the one hand, we should prove that $\omega \in A_p$ implies $\omega \in \mathrm{BMO}^{p'}(\omega)$.

Case 1: $p = 1$. Given a cube Q, it follows from the A_1 condition that

$$|\omega(x) - \omega_Q| \le \omega(x) + \omega_Q \le \big(1 + [\omega]_{A_1}\big)\omega(x),\ a.e.\ x \in Q,$$

which gives

$$\operatorname*{ess\,sup}_{x\in Q} \frac{|\omega(x) - \omega_Q|}{w(x)} \le 1 + [\omega]_{A_1}.$$

Case 2: $1 < p < \infty$. For any cube Q, we use Minkowskis inequality to establish

$$\begin{aligned}
&\left(\frac{1}{\omega(Q)}\int_Q |\omega(x) - \omega_Q|^{p'}\omega(x)^{1-p'}dx\right)^{1/p'} \\
&\le \frac{1}{\omega(Q)^{1/p'}}\left(\int_Q \omega(x)^{p'}\omega(x)^{1-p'}dx\right)^{1/p'} + \frac{\omega_Q}{\omega(Q)^{1/p'}}\left(\int_Q \omega(x)^{1-p'}dx\right)^{1/p'} \\
&:= I_1 + I_2.
\end{aligned}$$

It is easy to see that $I_1 = 1$. For I_2, $\omega \in A_p$ implies that

$$\begin{aligned}
I_2 &= \frac{1}{|Q|}\left(\int_Q \omega(x)dx\right)^{1-1/p'}\left(\int_Q \omega(x)^{1-p'}dx\right)^{1/p'} \\
&= \left(\frac{1}{|Q|}\int_Q \omega(x)dx\right)^{1/p}\left(\frac{1}{|Q|}\int_Q \omega(x)^{1-p'}dx\right)^{1/p'} \\
&\le [\omega]_{A_p}^{1/p}.
\end{aligned}$$

Therefore $\omega \in \mathrm{BMO}^{p'}(\omega)$ and $\|\omega\|_{\mathrm{BMO}^{p'}(\omega)} \le 1 + [\omega]_{A_p}^{1/p}$.

On the other hand, the condition $\omega \in A_p$ turns out to be necessary for the conclusion $\omega \in \mathrm{BMO}^{p'}(\omega)$ holds.

Case 1: $p = 1$. Given a cube Q, it follows from $\omega \in \mathrm{BMO}^{\infty}(\omega)$ that

$$\omega_Q \le |\omega(x) - \omega_Q| + \omega(x) \le \big(1 + \|\omega\|_{\mathrm{BMO}^{\infty}(\omega)}\big)\omega(x),\ a.e.\ x \in Q,$$

then, $M(\omega)(x) \le \big(1 + \|\omega\|_{\mathrm{BMO}^{\infty}(\omega)}\big)\omega(x), a.e. x \in \mathbb{R}^n$; that is, $\omega \in A_1$.

Case 2: $1 < p < \infty$. For any fixed cube Q, using Minkowski's inequality,

$$\begin{aligned}
\|\omega\|_{\mathrm{BMO}^{p'}(\omega)} &\ge \left(\frac{1}{w(Q)}\int_Q |\omega(y) - \omega_Q|^{p'}\omega(y)^{1-p'}dy\right)^{1/p'} \\
&\ge \left(\frac{1}{w(Q)}\int_Q \omega_Q^{p'}\omega(y)^{1-p'}dy\right)^{1/p'} - \left(\frac{1}{w(Q)}\int_Q \omega(y)^{p'}\omega(y)^{1-p'}dy\right)^{1/p'}
\end{aligned}$$

$$= \left(\frac{1}{|Q|}\int_Q \omega(x)dx\right)^{1/p}\left(\frac{1}{|Q|}\int_Q \omega(x)^{1-p'}dx\right)^{1/p'} - 1.$$

Thus, we conclude that $\omega \in A_p$ and $[\omega]_{A_p} \le \left(\|\omega\|_{\mathrm{BMO}^{p'}(\omega)} + 1\right)^p$.

Hence, it immediately follows that $\omega \in A_p$ if and only if $\omega \in \mathrm{BMO}^{p'}(\omega)$ with

$$[\omega]_{A_p}^{1/p} \le \|\omega\|_{\mathrm{BMO}^{p'}(\omega)} + 1 \le [\omega]_{A_p}^{1/p} + 2.$$

It is easy to see that for $1 \le p < \infty$, $[\omega]_{A_p} \ge 1$. Thus, the proof of Theorem 1.5 is completed. □

Proof of Theorem 1.7. For any $\omega \in A_\infty$, there exists $1 \le p_0 < \infty$ such that $\omega \in A_{p_0}$. Applying Theorem 1.6, we have

$$\omega \in \mathrm{BMO}^{p_0'}(\omega) \subset \mathrm{BMO}(\omega).$$

We finish the proof of Theorem 1.7. □

Remark 3.1. *For the case $p = \infty$. For any non-negative, locally integrable function ω,*

$$\|\omega\|_{\mathrm{BMO}(\omega)} = \sup_Q \frac{1}{\omega(Q)}\int_Q |\omega(x) - \omega_Q|dx \le 2.$$

Then $\omega \in A_\infty$ is not necessary for the conclusion $\omega \in \mathrm{BMO}(\omega)$ holds.

Indeed, the proof we just gave and Proposition 3.1 lead to the following result.

Theorem 3.2. *Let $1 \le p < \infty$, $\frac{1}{p} + \frac{1}{p'} = 1$ and $f \in L_{\mathrm{loc}}(\mathbb{R}^n)$. $|f| \in A_p$ if and only if $f \in \mathrm{BMO}^{p'}(|f|)$.*

Proof of Theorem 1.8. Since the case $p = \infty$ is trivial, it suffices to consider $1 \le p < \infty$. Let $f \in \mathrm{BMO}^p(\omega)$. For any cube Q, we use Fubini's theorem and Minkowski's inequality to establish

$$(U_\psi f)_Q = \int_0^1 \left(\frac{1}{|Q|}\int_Q f(tx)dx\right)\psi(t)dt = \int_0^1 f_{tQ}\psi(t)dt$$

and

$$\left(\frac{1}{w(Q)}\int_Q |U_\psi f(x) - (U_\psi f)_Q|^p \omega(x)^{1-p}dx\right)^{1/p}$$
$$\le \left(\frac{1}{w(Q)}\int_Q \left(\int_0^1 |f(tx) - f_{tQ}|\psi(t)dt\right)^p \omega(x)^{1-p}dx\right)^{1/p}$$

$$
\begin{aligned}
&\leq \int_0^1 \left(\frac{1}{\omega(Q)}\int_Q |f(tx)-f_{tQ}|^p\omega(x)^{1-p}dx\right)^{1/p}\psi(t)dt\\
&= \int_0^1 \frac{t^{(p-1)\alpha/p-n/p}}{\omega(Q)^{1/p}}\left(\int_{tQ} |f(x)-f_{tQ}|^p\omega(x)^{1-p}dx\right)^{1/p}\psi(t)dt\\
&\leq \|f\|_{\mathrm{BMO}^p(\omega)}\int_0^1 t^{(p-1)\alpha/p-n/p}\cdot\frac{\omega(tQ)^{1/p}}{\omega(Q)^{1/p}}\psi(t)dt\\
&= \|f\|_{\mathrm{BMO}^p(\omega)}\int_0^1 t^{(p-1)\alpha/p-n/p}\cdot\left(\frac{\int_{tQ}\omega(x)dx}{\int_Q\omega(x)dx}\right)^{1/p}\psi(t)dt\\
&= \|f\|_{\mathrm{BMO}^p(\omega)}\int_0^1 t^{\alpha}\psi(t)dt.
\end{aligned}
$$

Therefore, we have obtained the upper estimate

$$\|U_\psi f\|_{\mathrm{BMO}^p(\omega)\to\mathrm{BMO}^p(\omega)} \leq \int_0^1 t^\alpha\psi(t)dt.$$

To prove the opposite one, we take $f_0(x)=\omega(x)$. From Theorem 1.3 and the condition $\omega(tx)=t^\alpha\omega(x)$, it follows that $f_0(x)\in\mathrm{BMO}^p(\omega)$ and

$$U_\psi f_0(x) = \int_0^1 f_0(tx)\psi(t)dt = f_0(x)\int_0^1 t^\alpha\psi(t)dt.$$

If U_ψ is a bounded operator from $\mathrm{BMO}^p(\omega)$ to itself, then (1.2) holds. Meanwhile, if (1.2) holds, the constant $\int_0^1 t^\alpha\psi(t)dt$ is best possible. Thus we complete the proof. □

Given a nonnegative function ψ on $[0,1]$. For a measurable complex valued function f on $\mathbb{R}^n$, the weighted Cesàro average operator V_ψ, the adjoint operator of U_ψ, is defined by

$$(V_\psi f)(x) = \int_0^1 f(\frac{x}{t})t^{-n}\psi(t)dt.$$

The following result can be deduced immediately from the proof of Theorem 1.7.

Theorem 3.3. *Let $\psi:[0,1]\to[0,\infty)$ be a function, $p\in[1,\infty]$, $\alpha\in\mathbb{R}$, $\omega\in A_{p'}$ and $\omega(tx)=t^\alpha\omega(x)$ for all $t\in(0,\infty)$. Then, $V_\psi:\mathrm{BMO}^p(\omega)\to\mathrm{BMO}^p(\omega)$ exists as a bounded operator if and only if*

$$\int_0^1 t^{-\alpha-n}\psi(t)dt<\infty. \tag{2.3}$$

Moreover, when (2.3) holds, the operator norm of V_ψ on $\mathrm{BMO}^p(\omega)$ is given by

$$\|V_\psi\|_{\mathrm{BMO}^p(\omega)\to\mathrm{BMO}^p(\omega)} = \int_0^1 t^{-\alpha-n}\psi(t)dt.$$

4. Acknowledgement

I would like to thank the referee for the helpful comment which improved the presentation of this paper.

References

[1] Astala, K., Iwaniec,T., Saksman,E.: Beltrami operators in the plane, Duke Math. J. **107**(2001), 27-56.

[2] Buckley, S.: Estimates for operator norms on weighted spaces and reverse Jensen inequalities, Trans. Amer. Math. Soc. **340**(1993), 253-272.

[3] Bui, H., Paluszyński, M., Taibleson, M.: A maximal function characterization of weighted Besov-Lipschitz and Triebel-Lizorkin spaces, Studia Math. **119**(1996), 219-246 .

[4] Bui, H., Taibleson, M.: The characterization of the Triebel-Lizorkin space for $p = \infty$, J. Fourier Anal. Appl. **6**(2000), 537-550.

[5] Carton-Lebrun, C., Fosset, M.: Moyennes et quotients de Taylor dans BMO, Bull. Soc. Roy. Sci. Liége 53, **2**(1984), 85-87.

[6] Coifman, R., Fefferman, C.: Weighted norm inequalities for maximal functions and singular integrals, Studia Math. **51**(1974), 241-250.

[7] Dragičević, O., Grafakos, L., Pereyra, C. and Petermichl, S.: Extrapolation and sharp norm estimates for classical operators on weighted Lebesgue spaces, Publ. Math. **49**(2005), 73-91.

[8] García-Cuerva, J.: Weighted H^p spaces, Dissertationes Math. **162**(1979).

[9] Hunt, R., Muckenhoupt, B., Wheeden, R.: Weighted norm inequalities for the conjugate function and Hilbert transform, Trans. Amer. Math. Soc. **176**(1973), 227-251.

[10] Komori, Y., Shirai, S.: Weighted Morrey spaces and a singular integral operator, Math. Nachr. **282**(2009), 219-231.

[11] Lacey, M.T., Moen, K., Pérez, C., Torres, R.H.: Sharp weighted bounds for fractional integral operators, J. Funct. Anal. **259**(2010), 1073-1097.

[12] Martell, J.: Sharp maximal functions associated with approximations of the identity in spaces of homogeneous type and applications, Studia Math. **161**(2004), 113-145.

[13] Muckenhoupt, B.: Weighted norm inequalities for the Hardy maximal function, Trans. Amer. Math. Soc. **165**(1972), 207-226.

[14] Petermichl, S.: The sharp bound for the Hilbert transform in weighted Lebesgue spaces in terms of the classical A_p characteristic, Amer. J. Math. **129**(2007), 1355-1375.

[15] Petermichl, S.: The sharp weighted bound for the Riesz transforms, Proc. Amer. Math. Soc. **136**(2008), 1237-1249.

[16] Petermichl, S., Volberg, A.: Heating of the Ahlfors-Beurling operator: weakly quasiregular maps on the plane are quasiregular, Duke Math. J. **112**(2002), 281-305.

[17] Stein, E.M.: Harmonic analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals, Princeton University Press, (1993).

[18] Xiao, J.: L^p and BMO bounds of weighted Hardy-Littlewood averages, J Math Anal Appl. **262**(2001), 660-666.

College of Mathematics and System Sciences
Xinjiang University
Urumqi 830046
Republic of China
E-mail address: Wangdh1990@126.com; zhoujiangshuxue@126.com